



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.


We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**2734 [1918, 444]. Proposed by E. L. REES, University of Kentucky.**

Given two circles tangent to each other externally. From the extremity of a diameter through the point of tangency draw a secant such that the segment between the circles shall be equal to a given segment.

**SOLUTION BY THE PROPOSER.**

Let  $PB$  be the required position of the secant, assuming the problem having been solved. Draw  $PQ$  perpendicular to  $AC$  and let  $BP = y$ ,  $AQ = x$ ,  $AB = d$ ,  $AC = d'$ , and  $RP = a$ , the given segment. Then we have,  $y^2 = (d + x)^2 + x(d' - x)$ . Also  $(y - a)/d = (d + x)/y$ . Eliminating  $x$ , we get  $y^2 - 2a'y - d^2 = 0$ , where  $a' = a(2d + d')/2(d + d')$ . Hence,

$$y = a' \pm \sqrt{a'^2 + d^2}.$$

The analysis suggests the following construction: With  $B$  as a center,  $BC$  as a radius, describe the arc cutting the horizontal diameter produced of the lower circle in  $D$ .

Draw  $BD$  cutting the common tangent in  $E$ . Take  $EF = a$ . Draw  $FG$  perpendicular to  $AC$ . Then  $AG$  will equal  $a'$ . Make  $AH = AG$ . Draw  $BH$  and produce it to  $J$ , making  $HJ = AH$ . With  $B$  as a center and  $BJ$  as radius, describe an arc cutting the lower circle in  $P$ . Then  $BP$  is the required secant.

Also solved by P. J. DA CUNHA.

**2741 [1919, 35]. Proposed by H. L. OLSON, New Hampshire College.**

Prove or disprove the following statement: If the three sides and the area of a triangle are integers, at least one of the three altitudes is an integer.

**I. SOLUTION BY FRANK IRWIN, University of California.**

The statement is not true since the triangle whose sides are 5, 29, 30 has area 72, and altitudes  $144/5$ ,  $144/29$ , and  $24/5$ .

**II. REMARKS BY J. L. RILEY, Stephenville, Texas.**

On page 12 of Carmichael's *Diophantine Analysis* we find the following theorem:

A necessary and sufficient condition that rational numbers  $x, y, z$  shall represent the sides of a rational triangle is that they shall be proportional to numbers of the form  $n(m^2 + h^2)$ ,  $m(n^2 + h^2)$ ,  $(m + n)(mn - h^2)$ , where  $m, n, h$  are positive rational numbers and  $mn > h^2$ .

In deriving this result it is pointed out that if  $x = n(m^2 + h^2)$ ,  $y = m(n^2 + h^2)$ , and  $z = (m + n)(mn - h^2)$ , the area is  $hmn(m + n)(mn - h^2)$  and the altitude upon  $z$  is  $2hmn$ . If  $m, n$ , and  $h$  are integers ( $mn > h^2$ ) we have a series of triangles for which the statement of Mr. Olson is true.

**2742 [1919, 36]. Proposed by C. N. SCHMALL, New York City.**

In Gregory's *Examples in the Differential and Integral Calculus*, 1841, Chap. VII, p. 124, ex. 22, I find the following celebrated problem: "To find a point within a triangle from which if lines be drawn to the angular points their sum may be the least possible." The author remarks that "the direct solution of this problem is long and complicated, etc." Required a simple, brief solution.

**I. SOLUTION BY F. V. MORLEY, Johns Hopkins University.**

Consider the three points,  $\alpha, \beta, \gamma$  (complex variable) as on a unit or base circle. Then the sum of the distances from the point  $x$  is

$$(1) \quad \sum \sqrt{(x - \alpha) \left( \bar{x} - \frac{1}{\alpha} \right)}$$